

Tropospheric estimation using DORIS data: current status and perspectives

Pascal Willis (IGN/IPGP),


Yoaz Bar-Sever (JPL), Olivier Bock (IGN)

SUMMARY

- DORIS tropospheric estimation, where are we?
 - Current IGN strategy (ignwd08 for ITRF2008)
 - DORIS/GPS long term comparisons
 - DORIS/GPS/VLBI campaign (CONT08) comparisons
 - Estimating horizontal tropospheric gradients (comparisons + impact on station position)
- Post-ITRF2008 strategies

Current IGN analysis strategy


- A priori: function of height
- Mapping function: GMF --> VMF-1
- Parameter estimation:
 - Reset at start of pass
 - Only if time since last reset is larger than (20 minutes) --> not by pass but per station/time



DORIS/GPS long-term comparisons

(from Bock et al., Adv. Space Res., 2010)


2005-2008 DORIS/IGN vs GPS/IGS





GPS/DORIS tropo

GPS discontinuity

November 5, 2006 = 5 mm



Global comparisons (bias / standard deviations)




1

Possible problem related to SAA
(see also SPOT5 : [Stepanek et al., 2010](#))

CONT08 ZTD comparisons

(Teke et al., J. Geod., in press)

Koike Park, Hawaii




Ny Alesund, Norway




NB: much better results than for CONT02 (Snajdrova et al., J. Geod., 2006)

CONT08 - horizontal gradients



East




North

See also [Boehm et al., EGU 2010](#)

Horizontal tropospheric gradients (Willis et al., IAG Symp., in press)


- 2007 data : DORIS vs GPS/JPL/PPP




Goals:

- comparisons GPS/DORIS
- Impact on geodetic results


DORIS single station (Arequipa, daily results)



All stations




North
(best)




East

DORIS / GPS correlation



Mean value over 2007 (North)

North gradient / latitude



(slope predicted by McMillan et al., 1994)

Impact on station coordinates

Solution	Chi2/ DOF	North (mm)	East (mm)	Up (mm)
ignwd08	3.11	9.1	11.1	9.2
this study	2.39	8.8	11.5	9.0

TBD: reprocess all data (since 1993) and check repeatability

Post-ITRF2008 strategies (IGN)

- A priori: use GPT (TBC)
- Mapping function: VMF-1 (TBC)
- Elevation cutoff: 5 degree (TBC)
- Horizontal gradients (1 per day) (TBC)

NB: use of gradients needs to be coordinated between ACs (otherwise, it will introduce systematic effects)